



**PsyCheck**

Responding to mental health issues  
within alcohol and drug treatment

# **Program References**



# References

- American Journal of Psychiatry (2000).** Practice guideline for the treatment of patients with major depressive disorder (revision). American Psychiatric Association. *American Journal of Psychiatry*, 157(4 Suppl), 1–45.
- Andrews, G., Creamer, M., Crino, R., Hunt, C., Lampe, L., & Page, A. (2003).** *The Treatment of Anxiety Disorders: Clinician Guides and Patient Manuals (Second ed.)*. Melbourne: Cambridge University Press.
- Arendt, M., & Munk-Jorgenson, P. (2004).** Heavy cannabis users seeking treatment: Prevalence of psychiatric disorders. *Social Psychiatry and Psychiatric Epidemiology*, 39, 97–105.
- Ashendon, R., C. Silagy & Weller. (1997).** A systematic review of the effectiveness of promoting lifestyle change in general practice. *Family Practice* 14: 160-175.
- Ask, A., & Roche, A.M. (2005).** *Clinical Supervision: A Practical Guide for the Alcohol and Other Drugs Field*. Adelaide: National Centre for Education and Training on Addiction, Flinders University.
- Back, S. E., Dansky, B. S., Carroll, K. M., Foa, E. B., & Brady, K. T. (2001).** Exposure therapy in the treatment of PTSD among cocaine-dependent individuals: Description of procedures. *Journal of Substance Abuse Treatment*, 21(1), 35–45.
- Baillie, A. J., & Rapee, R. M. (2004).** Predicting who benefits from psychoeducation and self help for panic attacks. *Behaviour Research & Therapy*, 42(5), 513–527.
- Baker, A., & Wilson, P. H. (1985).** Cognitive therapy for depression: The effect of booster sessions on relapse. *Behaviour Therapy*, 6, 335–344.
- Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979).** *Cognitive therapy of depression*. New York: Guilford Press.
- Beck, J. S. (1995).** *Cognitive therapy: Basics and beyond*. New York: Guilford Press.
- Bernard, J. M. & Goodyear, R. K. (2004).** *Fundamentals of Clinical Supervision*. Boston: Pearson
- Bero, L. A., R. Grilli, J. M. Grimshaw, E. Harvey, A. D. Oxman, and M. A. Thomson. (1998).** Closing the Gap between Research and Practice: An Overview of Systematic Reviews of Interventions to Promote the Implementation of Research Findings. *British Medical Journal* 317 (7156): 465-8.
- Beusenberg M and Orley J. (1994)** *A user's guide to the Self Reporting Questionnaire (SRQ)*. Compiled by Division of Mental Health. Geneva: World Health Organization

## REFERENCES

- Bradley, A., & Toohey, B. (1999).** *The Coffs Harbour Project: A violence prevention program for substance misusing mentally ill.* Canberra: Commonwealth Department of Human Resources and Health, commissioned report.
- Brown, R. A., Evans, D. M., Miller, I. W., Burgess, E. S., & Mueller, T. I. (1997).** Cognitive-behavioral treatment for depression in alcoholism. *Journal of Consulting & Clinical Psychology*, 65(5), 715–726.
- Brown, S. A., Inaba, R. K., Gillin, J. C., Schuckit, M. A., Stewart, M. A., & Irwin, M. R. (1995).** Alcoholism and affective disorder: Clinical course of depressive symptoms. *American Journal of Psychiatry*, 152(1), 45–52.
- Burns, L., & Teesson, M. (2002).** Alcohol use disorders comorbid with anxiety, depression and drug use disorders. Findings from the Australian National Survey of Mental Health and Well Being. *Drug & Alcohol Dependence*, 68(3), 299–307.
- Callaly, T., Trauer, T., Munro, L., & Whelan, G. (2001).** Prevalence of psychiatric disorder in a methadone maintenance population. *Australian & New Zealand Journal of Psychiatry*, 35(5), 601–605.
- Carroll, K. M. (1998).** *Therapy Manuals for Drug Addiction : A Cognitive-Behavioral Approach: Treating Cocaine Addiction.* Maryland: National Institute on Drug Abuse.
- Craske, M. G., Brown, T. A., & Barlow, D. H. (1991).** Behavioural treatment of panic disorder: A two year follow-up. *Behavioural Therapy*, 22, 289–304.
- Crome, I. B., Christian, J., & Green, C. (2000).** The development of a unique designated community drug service for adolescents: Policy, prevention and education implications. *Drugs: Education, Prevention and Policy*, 7(1), 87–108.
- Darke, S., & Ross, J. (1997).** Polydrug dependence and psychiatric comorbidity among heroin injectors. *Drug & Alcohol Dependence*, 48(2), 135–141.
- Davis, D., Thomson, M.a., Freemantle, N., Wolf, F.m., Mazmanian, P.e., & Taylor-vaisey, A. (1999)** Impact of formal continuing medical education. Do conferences, workshops, rounds and other traditional continuing education activities change physician behavior or health care outcomes. *Journal of the American Medical Association*, 282, pp. 867-874.
- Dawe, S., Loxton, N. J., Hides, L., Kavanagh, D. J., & Mattick, R. P. (2002).** *Review of diagnostic and screening instruments for alcohol and other drug use and other psychiatric disorders (Second ed.).* Commonwealth of Australia, Publication Approval Number: 3146.
- Degenhardt, L., Hall, W., & Lynskey, M. (2001).** The relationship between cannabis use, depression and anxiety among Australian adults: Findings from the National Survey of Mental Health and Well-Being. *Social Psychiatry & Psychiatric Epidemiology*, 36(5), 219–227.
- DeRubeis, R. J., & Crits-Christoph, P. (1998).** Empirically supported individual and group psychological treatments for adult mental disorders. *Journal of Consulting & Clinical Psychology*, 66(1), 37–52.
- Egan, G. (1998).** *The skilled helper (Sixth ed.).* Pacific Grove, CA: Brooks/Cole.
- Ellis, A., & Harper, R. A. (1975).** *A new guide to rational living.* Englewood Cliffs, NJ: Wilshire.
- Ellis, P. M., Smith, D. A., & beyond blue: the national depression initiative (2002).** Treating depression: The beyondblue guidelines for treating depression in primary care. ‘Not so much what you do but that you keep doing it’. *Medical Journal of Australia*, 176, Suppl: S77–83.
- Feeney, S. (2003).** Seeking Safety: A treatment manual for PTSD and substance abuse. *Journal of Cognitive Psychotherapy*, 17, 293–295.

- Goldberg, D., & Williams, P. (1988).** *A User's Guide to the General Health Questionnaire*. Windsor, Berkshire, UK.: NFERNELSON Publishing Co. Ltd.
- Graham, H. L. (2004).** Implementing integrated treatment for coexisting substance use and severe mental health problems in assertive outreach teams: Training issues. *Drug and Alcohol Review*, 23, 463–470.
- Graham, H. L., Copello, A., Birchwood, M., Orford, J., McGovern, D., Atkinson, E., et al. (2000).** *Cognitive-behavioral integrated treatment (C-BIT): An approach for working with your clients who have severe mental health problems and use drugs/alcohol problematically*. Birmingham: Northern Birmingham Mental Health (NHS) Trust.
- Grant, B. F. (1997).** *The influence of comorbid major depression and substance use disorders on alcohol and drug treatment: Results of a national survey*. NIDA Research Monograph, 172. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism.
- Hall, W., & Farrell, M. (1997).** Comorbidity of mental disorders with substance misuse. *British Journal of Psychiatry*, 171, 4–5.
- Hart, G. M. (1982).** *The process of clinical supervision*. Baltimore: University Park Press.
- Heimberg, R. (2002).** Cognitive behavioural therapy for social anxiety disorder: Current status and future directions. *Biological Psychiatry*, 51, 101–108.
- Hollon, S. D., Haman, K. L., & Broan, L. L. (2002).** Cognitive-behavioural treatment of depression. In I. H. Gotlib & C. L. Hammen (Eds), *Handbook of Depression*, London: The Guilford Press.
- Hunter, S. B., Watkins, K. E., Wenzel, S., Gilmore, J., Shee, J., & Griffin, B. (2005).** Training substance abuse treatment staff to care for co-occurring disorders. *Journal of Substance Abuse Treatment*, 28, 239–245.
- Jarvis, T. J., Tebbutt, J., & Mattick, R. P. (2002).** *Treatment approaches for alcohol and drug dependence – An introductory guide*. Chichester: John Wiley.
- Kavanagh, D. J., Baker, A., & Teesson, M. (2004).** Special Section – Comorbidity of mental disorders and substance misuse: Introduction. *Drug and Alcohol Review*, 23(4), 405–406.
- Kavanagh, D. J., Greenaway, L., Jenner, L., Saunders, J. B., White, A., Sorban, J., Hamilton, G. (2000).** Contrasting views and experiences of health professionals on the management of comorbid substance misuse and mental disorders. *Australian and New Zealand Journal of Psychiatry*, 34(2), 279–289.
- Kay-Lambkin, F., Hazell, T. & Waring, T. (2000).** *PondROM: An educational resource on post-natal depression for general practitioners*. Newcastle, Australia: Hunter Institute of Mental Health.
- Kay-Lambkin, F. J., Baker, A. L., & Lewin, T. J. (2004).** The ‘co-morbidity roundabout’: A framework to guide assessment and intervention strategies and engineer change among people with comorbid problems. *Drug & Alcohol Review*, 23(4), 407–423.
- King, V. L., Stoller, K. B., Hayes, M., Umbricht, A., Currens, M., Kidorf, M. S., et al. (2002).** A multicenter randomized evaluation of methadone medical maintenance. *Drug and Alcohol Dependence*, 65(2), 137–148.
- Kirk, J. (1989).** *Cognitive-behavioral assessment*. In K. Hawton, P. M. Salkovskis, J. Kirk & D. M. Clark (Eds), *Cognitive behavior therapy for psychiatric problems: A practical guide*, 13–51. Oxford, UK: Oxford University Press.
- Krawitz, R., & Watson, C. (2003).** *Borderline personality disorder: a practical guide to treatment*. Oxford: Oxford University Press.

## REFERENCES

- Kushner, M. G., & Mueser, K. T. (1993).** *Psychiatric comorbidity with alcohol use disorders*. In Eighth Special Report to the US Congress on Alcohol and Health, NIH Publication No. 94-3699. Rockville: US Department of Health and Human Services.
- Ladouceur, R., Dugas, M. J., Freeston, M., Leger, E., Gagnon, F., & Thibodeau, N. (2000).** Efficacy of a cognitive-behavioural treatment for generalised anxiety disorder: Evaluation in a controlled clinical trial. *Journal of Consulting and Clinical Psychology*, 68(6), 957–964.
- Lee, N., Lintzeris, N., Ritter, A., & de Graaf, B. (2002).** *Psychosocial interventions for clients of methadone and buprenorphine maintenance: A systematic review and needs analysis*. Report to Department of Health and Aged Care. Canberra: Department of Health and Aged Care.
- Levesque, D. A., Prochaska, J. M., & Prochaska, J. O. (1999).** Stages of change and integrated service delivery. *Consulting Psychology Journal: Practice and Research*, 51(4), 226–241.
- Little, H.A., Rowe, C.L., Quille T.J, Dakof, G.A., Mills, D.S., Sakran E. & Biaggi, H. (2002)** Transporting a research-based adolescent drug treatment into practice. *Journal of Substance Abuse Treatment*, 22:4, pp231-243.
- McCloud, A., Barnaby, B., Omu, N., Drummond, C., & Aboud, A. (2004).** Relationship between alcohol use disorders and suicidality in a psychiatric population: In-patient prevalence study. *British Journal of Psychiatry*, 184, 439–445.
- Martin, G., Rozanes, P., Pearce, C., & Allison, S. (1995).** Adolescent suicide, depression and family dysfunction. *Acta Psychiatrica Scandinavica*, 92(5), 336-344.
- Maslin, J., Graham, H. L., Cawley, M., Copello, A., Birchwood, M., Georgiou, G., McGovern, D., Mueser, K. T. & Orford, J. (2001).** Combined severe mental health and substance use problems: What are the training and support needs of staff working with this client group? *Journal of Mental Health*, 19(2), 131–140.
- Minkoff, K. (2001).** Best practices: Developing standards of care for individuals with co-occurring psychiatric and substance use disorders. *Psychiatric Services*, 52, 597–599.
- NICS (2006).** *Taking action locally: Eight steps to putting cancer guidelines into practice*. Melbourne: National Institute of Clinical Studies.
- O’Leary, K. D., & Wilson, G. T. (1975).** *Behavior therapy: Application and outcome*. New Jersey: Prentice-Hall.
- Persons, J. B., Davidson, J., & Tompkins, M. A. (2001).** *Essential Components of Cognitive-Behaviour Therapy for Depression*. Washington: American Psychological Association.
- Powell D.J. & Brodsky, A (2004).** *Clinical Supervision in Alcohol and Drug Abuse Counseling: Principles, Models, Methods*. San Francisco: Jossey-Bass Inc
- Proudfoot, H., Teesson, M., Brewin, E., & Gournay, K. (2003).** *Comorbidity and delivery of services*. In H. Proudfoot (Ed.), *Comorbid Mental Disorders and Substance Use Disorders: Epidemiology, Prevention and Treatment*. Canberra: Commonwealth of Australia (publication approval number: 3415).
- Regier, D. A., Rae, D. S., Narrow, W. E., Kaelber, C. T., & Schatzberg, A. F. (1998).** Prevalence of anxiety disorders and their comorbidity with mood and addictive disorders. *British Journal of Psychiatry Supplementum*, 34, 24–28.
- Ridgely, M. S., Goldman, H. H., & Willenbring, M. (1990).** Barriers to the care of persons with dual diagnoses: Organizational and financing issues. *Schizophrenia Bulletin*, 116, 123–132.

- Roman, P. M., & Johnson, J. A. (2002).** Adoption and implementation of new technologies in substance abuse treatment. *Journal of Substance Abuse Treatment*, 22(4), 211–218.
- Schuckit, M. A., Tipp, J. E., Bergman, M., Reich, W., Hesselbrock, V. M., & Smith, T. L. (1997).** Comparison of induced and independent major depressive disorders in 2,945 alcoholics. *American Journal of Psychiatry*, 154, 948–957.
- Scogin, F. R., Hanson, A., & Welsh, D. (2003).** Self-administered treatment in stepped-care models of depression treatment. *Journal of Clinical Psychology*, 59(3), 341–349.
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002).** *Mindfulness-Based Cognitive Therapy for Depression: A New Approach to Preventing Relapse*. New York: The Guilford Press.
- Smith, S. S., Jorenby, D. E., Fiore, M. C., Anderson, J. E., Mielke, M. M., Beach, K. E., et al. (2001).** Strike while the iron is hot: Can stepped-care treatments resurrect relapsing smokers? *Journal of Consulting & Clinical Psychology*, 69(3), 429–439.
- Sobell, M. B., & Sobell, L. C. (2000).** Stepped care as a heuristic approach to the treatment of alcohol problems. *Journal of Consulting & Clinical Psychology*, 68(4), 573–579.
- Teesson, M., Burns, L., & National Drug and Alcohol Research Centre (2001).** *National Comorbidity Project*. Canberra: Commonwealth Department of Health and Aged Care.
- Teesson, M., Hall, W., Lynskey, M., & Degenhardt, L. (2000).** Alcohol and drug use disorders in Australia: Implications of the National Survey of Mental Health and Wellbeing. *Australian & New Zealand Journal of Psychiatry*, 34(2), 206–213.
- Van Noppen, B. L., Pato, M. T., Marsland, R., & Rasmussen, S. A. (1998).** A time-limited behavioral group for treatment of obsessive-compulsive disorder. *Journal of Psychotherapy Practice & Research*, 7(4), 272–280.
- Wallace, P., Cutler, S., & Haines, A. (1988).** Randomised controlled trial of general practitioner intervention in patients with excessive alcohol consumptions. *British Medical Journal*, 297(10), 663–668.
- Welch, M., & Mooney, J. (2001).** Managing services that manage people with a coexisting mental health and substance use disorder. *Australasian Psychiatry*, 9, 345–349.
- Westermeyer, J. J., Eames, S., & Nugent, S. (1998).** Comorbid dysthymia and substance use disorder: Treatment history and cost. *American Journal of Psychiatry*, 155, 1556–1560.
- Whitehead, L., & Royles, M. (2002).** Deliberate self-harm: assessment and treatment interventions. In S. Regel & D. Roberts (Eds.), *Mental health liaison: a handbook for nurses and health professionals*. Edinburgh, London: Baillière Tindall
- WHO (1997)** *Composite International Diagnostic Interview (CIDI) Version 2.1*. Geneva: World Health Organization (WHO)
- Williams, K. (1999).** Attitudes of mental health professionals to co-morbidity between mental health problems and substance misuse. *Journal of Mental Health*, 8, 605–613.
- Wilson, P. (1992).** *Principles and Practice of Relapse Prevention*. New York: The Guilford Press.

## REFERENCES